



ELSEVIER

Journal of Pure and Applied Algebra 105 (1995) 195–224

JOURNAL OF
PURE AND
APPLIED ALGEBRA

Group schemes and rigidity of algebras in positive characteristic

Serge Skryabin

Chebotaev Research Institute of Mathematics and Mechanics, Universitetskaya Str. 17, 420008 Kazan, Russia

Communicated by E.M. Friedlander; received 26 May 1993

Abstract

Rigidity criteria for a finite dimensional associative or Lie algebra of positive characteristic are given. A geometrically rigid algebra may have deformations with nontrivial infinitesimals which may be interpreted as obstructions to integrating infinitesimal automorphisms. A group scheme theoretic nature of those obstructions is revealed. For each affine group scheme G of finite type over the ground field an invariantly defined G -module $\text{Obs}(G)$ is introduced and formal properties of the functor $G \mapsto \text{Obs}(G)$ are studied.

1991 Math. Subj. Class.: 14D15, 14L15, 16S80

Introduction

Let L be a finite dimensional Lie algebra over a field k . Gerstenhaber and Schack [8, 9] introduced the terminology which distinguishes several concepts of rigidity. It has long been known that L is both analytically and geometrically rigid provided that the second cohomology group $H^2(L, L)$ vanishes [6, 13]. For an algebraically closed k this result can be improved as follows. Denote by Sq the canonical quadratic mapping $H^2(L, L) \rightarrow H^3(L, L)$ (see [16]). Let $\text{Ker Sq} = \{\zeta \in H^2(L, L) \mid \text{Sq } \zeta = 0\}$. Then L is both analytically and geometrically rigid provided that $\text{Ker Sq} = 0$. For geometric rigidity, one will find this result in a paper of Rauch [15, II, Theorem 3]. Further investigations showed that analytic rigidity implies geometric rigidity and that in the case of characteristic zero the converse is true [8]. Specific phenomena are encountered in characteristic $p > 0$. In this case $H^2(L, L)$ contains a subspace Obs which consists of obstructions to integrating the infinitesimal automorphisms. It was introduced by Gerstenhaber [7] in a more general context of composition complexes. We shall see that $\text{Sq } \zeta = 0$ for $\zeta \in \text{Obs}$. I do not know whether Sq factors through $H^2(L, L)/\text{Obs}$. Now we shall summarize the main results of the paper.